


Bovikalc Calcium Boluses For Cattle



Administration regime 1 Pre and post calving regime



First Bolus
Administer one bolus per animal
at the first sign of calving



Second Bolus
Follow up with second bolus
12-15 hours later if necessary

On-farm recommendation - if she's got cold ears, she needs another bolus

Administration regime 2 Post calving regime




First Bolus
Administer 1 bolus immediately after calving
(or after sub-cutaneous calcium injection)



Second Bolus
Follow up with a second bolus
12-15 hours later if necessary

On-farm recommendation - if she's got cold ears, she needs another bolus

Managing the calcium status of the herd during and after calving



Benefits

- Large supply of calcium - 43g in each bolus
- Bolus dissolves and supplies calcium within 20 minutes of administration
- Calcium from bolus available for 36 hours - peace of mind for the farmer
- None of the palatability problems associated with gels and pastes
- Can be used in conjunction with existing calcium supplementation regime
- **Active life - 36 hours**
- **Pack size - 4 boluses**



Aggrim Ltd The Flarepath, Elsham Wold Industrial Estate, Brigg, Lincolnshire, DN20 0SP, UK

Tel: +44 (0)1652 688046 Fax: +44 (0)1652 688049 Email: info@aggrim.co.uk Web: www.aggrim.co.uk

